

Invariant Relativistic Electrodynamics. Clifford Algebra Approach

Tomislav Ivezić

Ruđer Bošković Institute, P.O.B. 180, 10002 Zagreb, Croatia
ivezic@irb.hr

In the usual Clifford algebra formulation of electrodynamics the Faraday bivector field F is decomposed into *the observer dependent* sum of a relative vector $\mathbf{E}$ and a relative bivector $e_5\mathbf{B}$ by making a space-time split, which depends on the observer velocity. ($\mathbf{E}$ corresponds to the three-dimensional electric field vector, $\mathbf{B}$ corresponds to the three-dimensional magnetic field vector and e_5 is the (grade-4) pseudoscalar.) In this paper it is proved that the space-time split and the relative vectors are not relativistically correct, which means that the ordinary Maxwell equations with $\mathbf{E}$ and $\mathbf{B}$ and the field equations (FE) with F are not physically equivalent. Therefore we present *the observer independent* decomposition of F by using the 1-vectors of electric E and magnetic B fields. The equivalent, invariant, formulations of relativistic electrodynamics (independent of the reference frame and of the chosen coordination for that frame) which use F , E and B , the real multivector $\Psi = E - e_5cB$ and the complex 1-vector $\Psi = E - icB$ are developed and presented here. *The new observer independent FE* are presented in formulations with E and B , with real and complex Ψ . When the sources are absent the FE with real and complex Ψ become Dirac like relativistic wave equations for the free photon. The expressions for *the observer independent* stress-energy vector $T(v)$ (1-vector), energy density U (scalar), the Poynting vector S and the momentum density g (1-vectors), the angular momentum density M (bivector) and the Lorentz force K (1-vector) are directly derived from the FE. The local conservation laws are also directly derived from the FE and written in an invariant way.

*Henceforth space by itself, and time by itself, are doomed
to fade away into mere shadows and only a kind of union of
the two will preserve an independent reality. H. Minkowski*

I. INTRODUCTION

In the usual Clifford algebra treatments, e.g. [1 – 3], of electrodynamics the Maxwell equations (ME) are written as a single equation using the

electromagnetic field strength F (a bivector) and the gradient operator ∂ (1-vector). (As expressed in [3] (Found. Phys. **23**, 1295 (1993)) the reference [4] *Clifford Algebra to Geometric Calculus* is: "one of the most stimulating modern textbooks of applied mathematics, full of powerful formulas waiting for physical application.") In order to get the more familiar form the field bivector F is expressed in terms of the sum of a relative vector $\mathbf{E}$ (corresponds to the three-dimensional electric field vector) and a relative bivector $e_5\mathbf{B}$ ($\mathbf{B}$ corresponds to the three-dimensional magnetic field vector, and e_5 is the (grade-4) pseudoscalar) by making a space-time split, which depends on *the observer velocity*. It is considered in such formulation that the ME written in terms of F and of $\mathbf{E}$ and $\mathbf{B}$ are completely equivalent. The components of $\mathbf{E}$ and $\mathbf{B}$ are considered to define in a unique way the components of F . Moreover in order to get the wave theory of electromagnetism the vector potential A is introduced and F is defined in terms of A . Thus such formulation with relative vectors $\mathbf{E}$, $\mathbf{B}$ and with 1-vector A is not only observer dependent but also gauge dependent.

However in the recent works [5 – 7] an invariant formulation of special relativity (SR) is proposed (see also [8]) and compared with different experiments, e.g., the "muon" experiment, the Michelson-Morley type experiments, the Kennedy-Thorndike type experiments and the Ives-Stilwell type experiments. In such invariant formulation of SR a physical quantity in the 4D spacetime is mathematically represented either by a tensor (when no basis has been introduced) or equivalently by a coordinate-based geometric quantity (CBGQ) comprising both components and a basis (when some basis has been introduced). *This invariant formulation is independent of the reference frame and of the chosen coordination for that frame.* The CBGQs representing some 4D physical quantity in different relatively moving inertial frames of reference (IFRs), or in different coordinations of the chosen IFR, are all mathematically equal and thus they are *the same quantity* for different observers, or in different coordinations (this fact is the real cause for the name invariant SR). *It is taken in the invariant SR that such 4D tensor quantities are well-defined not only mathematically but also experimentally, as measurable quantities with real physical meaning.* The complete and well-defined measurement from this invariant SR viewpoint is such measurement in which all parts of some 4D quantity are measured. The invariant SR is compared with the usual covariant formulation, which mainly deals with the basis components of tensors in a specific, i.e., Einstein's coordination (EC). In the EC the Einstein synchronization [9] of distant clocks and Cartesian space coor-

ordinates x^i are used in the chosen IFR. Further the invariant SR is compared with the usual noncovariant approach to SR in which some quantities are not 4D tensor quantities, but rather quantities from "3+1" space and time, e.g., the synchronously determined spatial length (the Lorentz contraction) [9]. It is shown in [6] that all the experiments (when they are complete from the viewpoint of the invariant SR) are in agreement with that formulation but not always with the usual covariant or noncovariant approaches to SR. It is also found in [5] that the usual transformations of the 3D vectors of the electric and magnetic fields $\mathbf{E}$ and $\mathbf{B}$ are not relativistically correct.

In this paper it is shown that the space-time split is not relativistically correct procedure and that the relative vectors are not well-defined quantities from the SR viewpoint. This means that the ordinary ME with $\mathbf{E}$ and $\mathbf{B}$ are not physically equivalent with the observer independent FE with F . Further we write the Lorentz transformations (LT) in a coordination independent way. Then we present *the observer independent decomposition of F in terms of 1-vectors E and B* . The new Clifford algebra formulations of relativistic electrodynamics with 1-vectors E and B and with the real multivector $\Psi = E - e_5 c B$, or with the complex 1-vector $\Psi = E - icB$ (i is the unit imaginary), which are completely equivalent to the formulation with the field bivector F , are developed and presented here. The expressions for the observer independent stress-energy vector $T(v)$ (1-vector), energy density U (scalar, i.e., grade-0 multivector), the Poynting vector S (1-vector), the angular momentum density M (bivector) and the Lorentz force K (1-vector) are directly derived from the FE and given in all four formulations. Consequently *the principle of relativity is automatically included in such formulations with invariant quantities, whereas in the traditional formulation of SR this principle acts as the postulate established outside the mathematical formulation of the theory*. The local charge-current density and local energy-momentum conservation laws are derived from the FE. It is also shown that in the real and the complex Ψ formulations the FE become a Dirac like relativistic wave equation for the free photon. The expressions for such geometric 4D quantities are compared with the familiar ones from the 3D space considering our definitions in the standard basis $\{\gamma_\mu\}$ and in the $\mathcal{R}$ frame (the frame of "fiducial" observers) in which $E^0 = B^0 = 0$. This formalism does not make use of the intermediate electromagnetic 4-potential A , and thus dispenses with the need for the gauge conditions. The main idea for the whole approach is the same as for the invariant SR with tensors [5 – 8], i.e., that *the physical meaning is attributed, both theoretically and experimentally, only to the observer*

independent 4D quantities. We also remark that the observer independent quantities introduced here and the FE written in terms of them are of the same form both in the flat and curved spacetimes.

II. SHORT REVIEW OF GEOMETRIC ALGEBRA. SPACE-TIME SPLIT. LORENTZ TRANSFORMATIONS

A. A brief summary of geometric algebra

First we provide a brief summary of geometric algebra. We write Clifford vectors in lower case (a) and general multivectors (Clifford aggregate) in upper case (A). The space of multivectors is graded and multivectors containing elements of a single grade, r , are termed homogeneous and written A_r . The geometric (Clifford) product is written by simply juxtaposing multivectors AB . A basic operation on multivectors is the degree projection $\langle A \rangle_r$ which selects from the multivector A its r -vector part ($0 = \text{scalar}$, $1 = \text{vector}$, $2 = \text{bivector}$...). We write the scalar (grade-0) part simply as $\langle A \rangle$. The geometric product of a grade- r multivector A_r with a grade- s multivector B_s decomposes into $A_r B_s = \langle AB \rangle_{r+s} + \langle AB \rangle_{r+s-2} \dots + \langle AB \rangle_{|r-s|}$. The inner and outer (or exterior) products are the lowest-grade and the highest-grade terms respectively of the above series $A_r \cdot B_s \equiv \langle AB \rangle_{|r-s|}$, and $A_r \wedge B_s \equiv \langle AB \rangle_{r+s}$. For vectors a and b we have $ab = a \cdot b + a \wedge b$, where $a \cdot b \equiv (1/2)(ab + ba)$, and $a \wedge b \equiv (1/2)(\underline{ab} - \underline{ba})$. Reversion is an invariant kind of conjugation, which is defined by $\overline{AB} = \tilde{B}\tilde{A}$, $\tilde{a} = a$, for any vector a , and it reverses the order of vectors in any given expression. Also we shall need the operation called the complex reversion (for example, when working with complex 1-vector $\Psi = E - ciB$). The complex reversion of, e.g., Ψ , is denoted by an overbar $\overline{\Psi}$. It takes the complex conjugate of the scalar (complex) coefficient of each of the 16 elements in the algebra, and reverses the order of multiplication of vectors in each multivector.

B. Standard basis, non-standard bases, and the space-time split

In the treatments, e.g., [1 – 3], one usually introduces the standard basis. The generators of the spacetime algebra (STA) (the Clifford algebra generated by Minkowski spacetime) are taken to be four basis vectors $\{\gamma_\mu\}$, $\mu = 0, 1, 2, 3$, satisfying $\gamma_\mu \cdot \gamma_\nu = \eta_{\mu\nu} = \text{diag}(+ - - -)$. This basis is a right-handed orthonormal frame of vectors in the Minkowski spacetime M^4 with γ_0 in the forward light cone. The γ_k ($k = 1, 2, 3$) are spacelike vectors. This algebra is often called the Dirac algebra D and the elements of D are called

d -numbers. The γ_μ generate by multiplication a complete basis, the standard basis, for STA: $1, \gamma_\mu, \gamma_\mu \wedge \gamma_\nu, \gamma_\mu \gamma_5, \gamma_5$ ($2^4 = 16$ independent elements). γ_5 is the pseudoscalar for the frame $\{\gamma_\mu\}$.

We remark that the standard basis corresponds, in fact, to the specific coordination, i.e., the EC, of the chosen IFR. However different coordinations of an IFR are allowed and they are all equivalent in the description of physical phenomena. For example, in [5] two very different, but completely equivalent coordinations, the EC and "radio" ("r") coordination, are exposed and exploited throughout the paper. For more detail about the "r" coordination see, e.g., [5] and references therein.

The next step in the usual treatments, e.g., [1 – 3], is the introduction of a space-time split and the relative vectors. Since the usual STA deals exclusively with the EC it is possible to say that a given IFR is completely characterized by a single future-pointing, timelike unit vector γ_0 (γ_0 is tangent to the world line of an observer at rest in the γ_0 -system). By singling out a particular time-like direction γ_0 we can get a unique mapping of spacetime into the even subalgebra of STA (the Pauli subalgebra). For each spacetime point (or event) x this mapping is specified by

$$x\gamma_0 = ct + \mathbf{x}, \quad ct = x \cdot \gamma_0, \quad \mathbf{x} = x \wedge \gamma_0. \quad (1)$$

To each event x the equation (1) assigns a unique time and position in the γ_0 -system. The set of all position vectors $\mathbf{x}$ is the 3-dimensional position space of the observer γ_0 and it is designated by $P^3 = P^3(\gamma_0) = \{\mathbf{x} = x \wedge \gamma_0\}$. The elements of P^3 are all spacetime bivectors with γ_0 as a common factor ($x \wedge \gamma_0$). They are called *the relative vectors* (relative to γ_0) and they will be designated in boldface. Then a standard basis $\{\sigma_k; k = 1, 2, 3\}$ for P^3 , which corresponds to a standard basis $\{\gamma_\mu\}$ for M^4 is given as $\sigma_k = \gamma_k \wedge \gamma_0 = \gamma_k \gamma_0$. The invariant distance x^2 then decomposes as $x^2 = (x\gamma_0)(\gamma_0 x) = (ct - \mathbf{x})(ct + \mathbf{x}) = c^2 t^2 - \mathbf{x}^2$. The explicit appearance of γ_0 in (1) imply that *the space-time split is observer dependent*, i.e., it is dependent on the chosen IFR. It has to be noted that in the EC the space-time split of the position 1-vector x (1) gives separately the space and time components of x with their usual meaning, i.e., as in the prerelativistic physics, and (as shown above) in the invariant distance x^2 the spatial and temporal parts are also separated. (In the "r" coordination there is no space-time split and also in x^2 the spatial and temporal parts are not separated, see [5].) This does not mean that the EC does have some advantage relative to other coordinations and that the quantities in the EC are more physical than, e.g., those in the "r" coordination.

Different coordinations refer to the same IFR, say the S frame. But if we consider the geometric quantity, the position 1-vector, x in another relatively moving IFR S' , which is characterized by γ'_0 , then the space-time split in S' and in the EC is $x\gamma'_0 = ct' + \mathbf{x}'$, and this $x\gamma'_0$ is not obtained by the LT (or any other coordinate transformations) from $x\gamma_0$. (The hypersurface $t' = \text{const.}$ is not connected in any way with the hypersurface $t = \text{const.}$) Thus the customary Clifford algebra approaches to SR start with the geometric, i.e., coordinate-free, quantities, e.g., x, x^2 , etc., which are physically well-defined. However the use of the space-time split introduces in the customary approaches such coordinate-dependent quantities which are not physically well-defined since they cannot be connected by the LT. The main difference between our invariant approach to SR (by the use of the Clifford algebra) and the other Clifford algebra approaches is that in our approach, as already said, *the physical meaning is attributed, both theoretically and experimentally, only to the geometric 4D quantities, and not to their parts. Thus there is no need and moreover it is not physical from the viewpoint of the invariant SR to introduce the space-time split of the geometric 4D quantity.* We consider, in the same way as H. Minkowski (the motto in this paper), that the spatial and the temporal components (e.g., $\mathbf{x}$ and t , respectively) of some geometric 4D quantity (e.g., x) are not physically well-defined quantities. Only their union is physically well-defined and only such quantity does have an independent reality.

Thus instead of the standard basis $\{\gamma_\mu\}$, $\mu = 0...3$, for M^4 we can use some basis $\{e_\mu\}$ (the metric tensor of M^4 is then defined as $g_{\mu\nu} = e_\mu \cdot e_\nu$) and its dual basis $\{e^\mu\}$, where the set of base vectors e^μ is related to the e_μ by the conditions $e_\mu \cdot e^\nu = \delta_\mu^\nu$. The pseudoscalar e_5 of a frame $\{e_\mu\}$ is defined by $e_5 = e_0 \wedge e_1 \wedge e_2 \wedge e_3$. Then, e.g., the position 1-vector x can be decomposed in the S and S' frames and in the standard basis $\{\gamma_\mu\}$ and some non-standard basis $\{e_\mu\}$ as $x = x^\mu \gamma_\mu = x^{\mu'} \gamma_{\mu'} = \dots = x_e^{\mu'} e_{\mu'}$. The primed quantities are the Lorentz transforms of the unprimed ones. Similarly any multivector A can be written as an invariant quantity with the components and the basis, i.e., as the CBGQ. In such interpretation the LT are considered as passive transformations; both the components and the base vectors are transformed but the whole geometric quantity remains unchanged. Thus we see that under the passive LT a well-defined quantity on the 4D spacetime, i.e., a CBGQ, is an invariant quantity. This doesn't hold for the relative vectors and thence they are not well-defined quantities from the SR viewpoint.

C. Lorentz transformations

In the usual Clifford algebra formalism, e.g., [1–4], the LT are considered as active transformations; the components of, e.g., some 1-vector relative to a given IFR (with the standard basis $\{\gamma_\mu\}$) are transformed into the components of a new 1-vector relative to the same frame (the basis $\{\gamma_\mu\}$ is not changed). Furthermore the LT are described with rotors R , $R\tilde{R} = 1$, in the usual way as $p \rightarrow p' = Rp\tilde{R} = p_{\mu'}\gamma^{\mu'}$. But every rotor in spacetime can be written in terms of a bivector as $R = e^{\theta/2}$. For boosts in arbitrary direction $e^{\theta/2} = (1 + \gamma + \gamma\beta\gamma_0 n)/(2(1 + \gamma))^{1/2}$, $\theta = \alpha\gamma_0 n$, β is the scalar velocity in units of c , $\gamma = (1 - \beta^2)^{-1/2}$, or in terms of an ‘angle’ α we have $\tanh \alpha = \beta$, $\cosh \alpha = \gamma$, $\sinh \alpha = \beta\gamma$, and n is not the basis vector but any unit space-like vector orthogonal to γ_0 ; $e^\theta = \cosh \alpha + \gamma_0 n \sinh \alpha$. (One can also express the relationship between the two relatively moving frames S and S' in terms of rotor as $\gamma_{\mu'} = R\gamma_\mu\tilde{R}$.) The above explicit form for $R = e^{\theta/2}$ is frame independent but it is coordination dependent since it refers to the EC.

Here a coordination independent form for the LT is introduced and it can be used both in an active way (when there is no basis) or in a passive way (when some basis is introduced). The main step is the introduction of the 1-vector $u = cn$, which represents *the proper velocity of the frame S with respect to itself*. Then taking that v is 1-vector of the velocity of S' relative to S we write the component form of L in some basis $\{e_\mu\}$ which, as already said, does not need to be the standard basis, as

$$L_\nu^\mu = g_\nu^\mu + 2u^\mu v_\nu c^{-2} - (u^\mu + v^\mu)(u_\nu + v_\nu)/c^2(1 + u \cdot v/c^2), \quad (2)$$

or with the components and the basis, i.e., as the CBGQ, $L = L_\nu^\mu e_\mu e^\nu$, see the second paper in [8] and [5]; actually this form of the LT is a generalization to arbitrary coordination of the covariant form of the LT in the EC given in [10]. The rotor connected with such L is

$$R = L/(\tilde{L}L)^{1/2} = L_\nu^\mu e_\mu e^\nu/(\tilde{L}L)^{1/2}, \quad \tilde{L}L = 8(\gamma + 1), \quad \gamma = u \cdot v/c^2, \quad (3)$$

It can be also written as

$$\begin{aligned} R &= \langle R \rangle + \langle R \rangle_2 = \cosh \alpha/2 + ((u \wedge v)/|u \wedge v|) \sinh \alpha/2 = \\ &\quad \exp((\alpha/2)(u \wedge v)/|u \wedge v|), \\ R &= ((1 + u \cdot v)/2)^{1/2} + ((u \wedge v)/|u \wedge v|)((-1 + u \cdot v)/2)^{1/2}. \end{aligned} \quad (4)$$

One also can solve L_ν^μ in terms of R as

$$L_\nu^\mu = \langle e_\nu \tilde{R} e^\mu R \rangle. \quad (5)$$

The usual results are recovered when the standard basis $\{\gamma_\mu\}$, i.e., the EC is used. But these results for L and R hold also for other bases, i.e., coordinations. (Thus one can easily find the LT in the "r" coordination, $L_{\nu,r}^\mu$, and compare it with the corresponding result in [5].)

III. THE F FORMULATION OF ELECTRODYNAMICS AND THE PROOF THAT THE SPACE-TIME SPLIT AND THE TRANSFORMATIONS OF RELATIVE VECTORS $\mathbf{E}$ AND $\mathbf{B}$ ARE NOT RELATIVISTICALLY CORRECT

A. The F formulation of electrodynamics

We start the exposition of electrodynamics writing the FE in terms of F [1 – 3]; an electromagnetic field is represented by a bivector-valued function $F = F(x)$ on spacetime. The source of the field is the electromagnetic current j which is a 1-vector field. Then using that the gradient operator ∂ is a 1-vector FE can be written as a single equation

$$\partial F = j/\varepsilon_0 c, \quad \partial \cdot F + \partial \wedge F = j/\varepsilon_0 c. \quad (6)$$

The trivector part is identically zero in the absence of magnetic charge. Notice that in [1 – 3] the FE (6) are considered to encode all of the ME, i.e., that the FE (6) and the usual ME with $\mathbf{E}$ and $\mathbf{B}$ are physically equivalent. Our discussion will show that this is not true.

The field bivector F yields the complete description of the electromagnetic field and, in fact, there is no need to introduce either the field vectors or the potentials. For the given sources the Clifford algebra formalism enables one to find in a simple way the electromagnetic field F . Namely the gradient operator ∂ is invertible and (6) can be solved for $F = \partial^{-1}(j/\varepsilon_0 c)$, see, e.g., [1 – 3].

In the Clifford algebra formalism one can easily derive the expressions for the stress-energy vector $T(v)$ and the Lorentz force K directly from FE (6) and from the equation for $\tilde{F}$, the reverse of F , $\tilde{F}\tilde{\partial} = \tilde{j}/\varepsilon_0 c$ ($\tilde{\partial}$ differentiates to the left instead of to the right). Indeed, using (6) and from the equation for $\tilde{F}$ one finds

$$T(\partial) = (-\varepsilon_0/2)(F\partial F) = j \cdot F/c = -K, \quad (7)$$

where in $(F\partial F)$ the derivative ∂ operates to the left and to the right by the chain rule. The stress-energy vector $T(v)$ [1 – 3] for the electromagnetic field is then defined in the F formulation as

$$T(v) = T(v(x), x) = -(\varepsilon_0/2c) \langle FvF \rangle_1. \quad (8)$$

We note that $T(v)$ is a vector-valued linear function on the tangent space at each spacetime point x describing the flow of energy-momentum through a surface with normal $n = n(x)$; $v = cn$.

The right hand side of (7) yields the expression for the Lorentz force K , $K = F \cdot j/c$. This relation shows that the Lorentz force K can be interpreted as the rate of energy-momentum transfer from the source j to the field F . The Lorentz force in the F formulation for a charge q is $K = (q/c)F \cdot u$, where u is the velocity 1-vector of a charge q (it is defined to be the tangent to its world line).

The stress-energy vector $T(v)$ can be written in the following form

$$T(v) = -(\varepsilon_0/2c) [(F \cdot F)v + 2(F \cdot v) \cdot F]. \quad (9)$$

We write $T(v)$ (9) in a *new form* as a sum of v -parallel part ($v - \parallel$) and v -orthogonal part ($v - \perp$)

$$\begin{aligned} T(v) = & -(\varepsilon_0/2c) \left[-(F \cdot F) + (2/c^2)(F \wedge v)^2 \right] v + \\ & -(\varepsilon_0/c) \left[(F \cdot v) \cdot F - (1/c^2)(F \cdot v)^2 v \right]. \end{aligned} \quad (10)$$

The first term in (10) is $v - \parallel$ part and it yields the energy density U . Namely using $T(v)$ and the fact that $v \cdot T(v)$ is positive for any timelike vector v we construct the expression for *the observer independent energy density* U contained in an electromagnetic field as $U = v \cdot T(v)/c = (1/c) \langle vT(v) \rangle$, (scalar, i.e., grade-0 multivector). Thus in terms of F and (10) U becomes

$$U = (-\varepsilon_0/2c^2) \langle FvFv \rangle = (\varepsilon_0/2) \left[(F \cdot F) - (2/c^2)(F \wedge v)^2 \right]. \quad (11)$$

The second term in (10) is $v - \perp$ part and it is $(1/c)S$, where S is *the observer independent expression for the Poynting vector* (1-vector),

$$S = -\varepsilon_0 \left[(F \cdot v) \cdot F - (1/c^2)(F \cdot v)^2 v \right], \quad (12)$$

and, as can be seen, $v \cdot S = 0$. Thus $T(v)$ expressed by U and S is

$$T(v) = (1/c)(Uv + S). \quad (13)$$

The observer independent momentum density g is defined as $g = (1/c^2)S$, i.e., g is $(1/c)$ of the v - $\perp$ part from (10)

$$g = -(\varepsilon_0/c^2) \left[(F \cdot v) \cdot F - (1/c^2)(F \cdot v)^2 v \right]. \quad (14)$$

From $T(v)$ (10) one finds also the expression for the observer independent angular-momentum density M

$$M = (1/c)T(v) \wedge x = (U/c^2)v \wedge x + g \wedge x. \quad (15)$$

The first term in (15) corresponds to the "orbital" angular-momentum density whereas the second term yields the "spin" or intrinsic angular-momentum density. It has to be emphasized once again that all these definitions are the definitions of invariant quantities, i.e., frame and coordination independent quantities.

All these quantities can be written in some basis $\{e_\mu\}$, which does not need to be the standard basis, as CBGQs. Thus $T(v)$ (9) becomes

$$T(v) = -(\varepsilon_0/2c) \left[(-1/2)F^{\alpha\beta}F_{\alpha\beta}v^\rho e_\rho + 2F^{\alpha\beta}F_{\alpha\rho}v^\rho e_\beta \right], \quad (16)$$

the energy density U (11) is

$$U = (\varepsilon_0/2) \left[(1/2)F^{\alpha\beta}F_{\alpha\beta} - (2/c^2)F^{\alpha\beta}F_{\alpha\rho}v^\rho v_\beta \right], \quad (17)$$

and the Poynting vector S (12) becomes

$$S = -\varepsilon_0 \left[F^{\alpha\beta}F_{\alpha\rho}v^\rho e_\beta - (1/c^2)F^{\alpha\beta}F_{\alpha\rho}v^\rho v_\beta v^\lambda e_\lambda \right]. \quad (18)$$

(The energy-momentum tensor $T^{\mu\nu}$ in the F (and the E, B) formulations will be presented in Sec. IV C.)

B. The local conservation laws in the F - formulation

It is well-known that from the FE in the F - formulation (6) one can derive a set of conserved currents. Of course the same holds for all other formulations; the E, B -formulation, the real and the complex Ψ -formulation, which will be discussed below. Thus, for example, in the F - formulation one

derives in the standard way that j from (6) is a conserved current. Simply, the vector derivative ∂ is applied to the FE (6) which yields

$$(1/\varepsilon_0 c)\partial \cdot j = \partial \cdot (\partial \cdot F).$$

Using the identity $\partial \cdot (\partial \cdot M(x)) \equiv 0$ ($M(x)$ is a multivector field) one obtains *the local charge conservation law*

$$\partial \cdot j = 0. \quad (19)$$

In a like manner we find from (7) that

$$\partial \cdot T(v) = 0 \quad (20)$$

for the free fields. This is a *local energy-momentum conservation law*. In the derivation of (7) we used the fact that $T(v)$ is symmetric, i.e., that $a \cdot T(b) = T(a) \cdot b$. Namely using overdots the expression for $T(\partial)$ ($T(\partial) = (-\varepsilon_0/2)(F\partial F)$, where ∂ operates to the left and to the right by the chain rule) can be written as $T(\partial) = \dot{T}(\dot{\partial}) = (-\varepsilon_0/2)(\dot{F}\dot{\partial}F + F\dot{\partial}\dot{F}) = 0$, since in the absence of sources $\partial F = \dot{F}\dot{\partial} = 0$ (the overdot denotes the multivector on which the derivative acts). Then from the above mentioned symmetry of T one finds that $\dot{T}(\dot{\partial}) \cdot v = \partial \cdot T(v) = 0$, $\forall$ *const.* v , which proves the equation (20).

Inserting the expression (13) for $T(v)$ into the local energy-momentum conservation law (20) we find

$$(v \cdot \partial)U + \partial \cdot S = 0. \quad (21)$$

The relation (21) is the well-known Poynting's theorem but now completely written in terms of the observer independent quantities. Let us introduce the standard basis $\{\gamma_\mu\}$, i.e., an IFR with the EC, and in the $\{\gamma_\mu\}$ basis we choose that $v = c\gamma_0$, or in the component form it is $v^\mu(c, 0, 0, 0)$. Then the familiar form of Poynting's theorem is recovered in such coordinate system

$$\partial U / \partial t + \partial_i S^i = 0, \quad i = 1, 2, 3. \quad (22)$$

It is worthwhile to note that although U (11) and S (10), taken separately, are well-defined observer independent quantities, the relations (13), (20) and (21) reveal that only $T(v)$ (13), as a whole quantity, i.e., the combination of

U and S , enters into a fundamental physical law, the local energy-momentum conservation law (20). Thence one can say that only $T(v)$ (13), as a whole quantity, does have a real physical meaning, or, better to say, a physically correct interpretation. An interesting example that emphasizes this point is the case of a uniformly accelerated charge. In the usual (3D) approach to the electrodynamics ([11] Sec. 6.8.) the Poynting vector S is interpreted as an energy flux due to the propagation of fields. In such an interpretation it is not clear how the fields propagate along the axis of motion since for the field points on the axis of motion one finds that $S = 0$ (there is no energy flow) but at the same time $U \neq 0$ (there is an energy density). Our approach reveals that the important quantity is $T(v)$ and not S and U taken separately. $T(v)$ is $\neq 0$ everywhere on the axis of motion and the local energy-momentum conservation law (20) holds everywhere.

Of course the same law (20) will be obtained in other formulations, E and B , real and complex Ψ formulations, as well. In the same way one can derive the local angular momentum conservation law, see [1], Space-Time Calculus.

C. The space-time split and the relative vectors $\mathbf{E}$ and $\mathbf{B}$

As already said in the usual Clifford algebra treatments of the electromagnetism the field bivector F is expressed in terms of the sum of a relative vector $\mathbf{E}$ and a relative bivector $\gamma_5 \mathbf{B}$ by making a space-time split in the γ_0 frame

$$\begin{aligned} F &= \mathbf{E} + c\gamma_5 \mathbf{B}, \quad \mathbf{E} = (F \cdot \gamma_0)\gamma_0 = (1/2)(F - \gamma_0 F \gamma_0), \\ \gamma_5 \mathbf{B} &= (F \wedge \gamma_0)\gamma_0 = (1/2c)(F + \gamma_0 F \gamma_0). \end{aligned} \quad (23)$$

F can be written as the CBGQ in the standard basis $\{\gamma_\mu\}$ as

$$F = (1/2)F^{\mu\nu}\gamma_\mu \wedge \gamma_\nu = F^{0k}\gamma_0 \wedge \gamma_k + (1/2)F^{kl}\gamma_k \wedge \gamma_l, \quad k, l = 1, 2, 3. \quad (24)$$

From (24) and (23) one concludes that the relative vectors $\mathbf{E}$ and $\mathbf{B}$ are expressed in the standard basis $\{\gamma_\mu\}$ as

$$\mathbf{E} = F^{0k}\gamma_0 \wedge \gamma_k, \quad \gamma_5 \mathbf{B} = (1/2c)F^{kl}\gamma_k \wedge \gamma_l. \quad (25)$$

We see from (24) and (25) that the components of F in the $\{\gamma_\mu\}$ basis give rise to the tensor $F^{\mu\nu} = \gamma^\nu \cdot (\gamma^\mu \cdot F) = (\gamma^\nu \wedge \gamma^\mu) \cdot F$, which, written out as a matrix, has entries $E_i = -F^{0i}$ and $B_i = -(1/2)\varepsilon_{ijk}F^{jk}$. (We write the

components E_i and B_i (and ε_{ijk}) with lowered (generic) subscripts, since they are not the spatial components of the well-defined quantities on the 4D spacetime.) It is considered in such formulation that the FE written in terms of F and the ME with $\mathbf{E}$ and $\mathbf{B}$ (taken in the $\{\gamma_\mu\}$ basis) are completely equivalent. Such usual interpretation is physically meaningless since F is the well-defined geometric quantity in the 4D spacetime with the correct transformation properties while it is not the case for $\mathbf{E}$ and $\mathbf{B}$, as will be shown below. We consider, in accordance with Minkowski's assertion (the motto here), that physically meaningful is only the whole geometric quantity, e.g., the field bivector F , the position 1-vector x , etc. (when there is no basis) or the corresponding CBGQ $(1/2)F^{\mu\nu}e_\mu \wedge e_\nu$, $x^\mu e_\mu$, (when some basis $\{e_\mu\}$ is chosen), and not some parts of it, taken in the specific representation, e.g., ct and $\mathbf{x}$, or $\mathbf{E}$ and $\mathbf{B}$, etc. Such parts of a geometric quantity have no definite physical sense since they do not transform properly under the LT. Namely the active LT transform *all components together* leaving the basis unchanged, while the passive LT transform both, *all components and all basis vectors together* leaving the whole quantity unchanged. Hence the space-time split (in the EC) of any geometric quantity defined on the 4D spacetime is, in fact, an incorrect procedure from the point of view of SR. Also it cannot be said, as usually argued both in the tensor formalism and in the Clifford algebra formalism, that $F^{\mu\nu}$ (and thus F too) are determined by the components of 3D quantities $\mathbf{E}$ and $\mathbf{B}$. F is the geometric 4D quantity and for the given sources it is completely determined in an observer independent way by the equation $F = \partial^{-1}(j/\varepsilon_0 c)$. We repeat once again that *in the 4D spacetime the physical meaning is attributed only to the whole 4D geometric quantity not to its spatial and temporal parts*. This is the main difference between our approach and others, e.g., [1 – 3].

D. The proof that the transformations of relative vectors $\mathbf{E}$ and $\mathbf{B}$ are not relativistically correct

Let us now explicitly show that the above decomposition of F (23) is not relativistically correct and that the usual transformations of $\mathbf{E}$ and $\mathbf{B}$ are not the LT of quantities that are well-defined on the 4D spacetime. It can be easily seen that the LT (the active ones) of the field bivector F , $F' = RF\tilde{R}$, with $\theta = \alpha\gamma_0\gamma_1$ (all in the standard basis $\{\gamma_\mu\}$), yields

$$\begin{aligned}
F' = & ((1 + \gamma)/2)[F^{0k}\gamma_0 \wedge \gamma_k + (1/2)F^{kl}\gamma_k \wedge \gamma_l] + \gamma\beta F^{0k}\gamma_k \wedge \gamma_1 + \\
& \gamma\beta\gamma_0[(1/2)F^{kl}\gamma_k \wedge \gamma_l - (1/2)F^{kl}\gamma_k \wedge \gamma_l\gamma_1] -
\end{aligned} \tag{26}$$

$$(\gamma^2\beta^2/2(1+\gamma))[\gamma_0\gamma_1F^{0k}\gamma_0\wedge\gamma_k\gamma_0\gamma_1+\gamma_0\gamma_1(1/2)F^{kl}\gamma_k\wedge\gamma_l\gamma_0\gamma_1].$$

Using (25) one can write (26) as

$$\begin{aligned} F' = & ((1+\gamma)/2)(\mathbf{E}+c\gamma_5\mathbf{B})+\gamma\beta F^{0k}\gamma_k\wedge\gamma_1+\gamma\beta\gamma_0(\gamma_1\cdot c\gamma_5\mathbf{B})- \\ & ((1-\gamma)/2)\gamma_0\gamma_1(\mathbf{E}+c\gamma_5\mathbf{B})\gamma_0\gamma_1. \end{aligned} \quad (27)$$

The relations (26) and (27) clearly show that *the active LT do not transform F (24) into $F^{0'k'}\gamma_0\wedge\gamma_k+(1/2)F^{k'l'}\gamma_k\wedge\gamma_l$ but introduce some additional terms (e.g., $\gamma\beta F^{0k}\gamma_k\wedge\gamma_1$). The separated parts ($\mathbf{E}$ and $\mathbf{B}$) of a well-defined 4D quantity F do not transform into the corresponding transformed parts ($F^{0'k'}\gamma_0\wedge\gamma_k$ and $(1/2)F^{k'l'}\gamma_k\wedge\gamma_l$) Such result implies that the relative vectors $\mathbf{E}$ and $\mathbf{B}$ are not well-defined quantities on the 4D spacetime, since they do not have the correct transformation properties. Hence it is not possible to write, as usually supposed, that the transformed F' in the γ_0 frame does have the same form as F , i.e., that $F' = \mathbf{E}' + c\gamma_5\mathbf{B}'$, where it is interpreted that the LT of $\mathbf{E}$ and $\mathbf{B}$ are*

$$\begin{aligned} \mathbf{E}' &= (1+\gamma)/2\mathbf{E}+\gamma\beta\gamma_0(\gamma_1\cdot c\gamma_5\mathbf{B})+((\gamma-1)/2)\gamma_0\gamma_1\mathbf{E}\gamma_0\gamma_1, \\ c\gamma_5\mathbf{B}' &= ((1+\gamma)/2)c\gamma_5\mathbf{B}+\gamma\beta F^{0k}\gamma_k\wedge\gamma_1+ \\ & ((\gamma-1)/2)\gamma_0\gamma_1(c\gamma_5\mathbf{B})\gamma_0\gamma_1. \end{aligned} \quad (28)$$

The relations (28) are completely meaningless from the SR viewpoint and they have nothing to do with the LT of a 4D quantity. In general, the LT of some parts of a well-defined 4D quantity are not mathematically correct. Hestenes in "New Foundations for Classical Mechanics," [3] p.625, declares: "Considering the simplicity of the transformation law (3.48) (our $F' = RF\tilde{R}$) for F , it is obviously preferable to treat the electromagnetic field $F = \mathbf{E} + c\gamma_5\mathbf{B}$ as a unit, rather than transform $\mathbf{E}$ and $\mathbf{B}$ separately by (3.51a,b) (our (28))." Our objection is that this is not the question of the preferability but the question of the correctness. The transformations (28) are not less preferable than the transformation $F' = RF\tilde{R}$ but the transformations (28) are, as we said, mathematically (and physically) incorrect. Thence the same holds for the decomposition $F = \mathbf{E} + c\gamma_5\mathbf{B}$ and, more generally, for the space-time split of any well-defined 4D physical quantity.

The similar result can be obtained with the passive LT. The passive LT transform always the whole 4D quantity, basis and components, leaving the whole quantity unchanged. This does not hold if F is decomposed into the

relative vectors $\mathbf{E}$ and $\mathbf{B}$. Namely under the passive LT it must hold that $F = (1/2)F^{\mu\nu}\gamma_\mu \wedge \gamma_\nu = (1/2)F^{\mu'\nu'}\gamma_{\mu'} \wedge \gamma_{\nu'}$ (the primed quantities are the Lorentz transforms of the unprimed ones), which will not be fulfilled if F is written in terms of $\mathbf{E}$ and $\mathbf{B}$, i.e., as $F = \mathbf{E} + c\gamma_5\mathbf{B}$. The relative vectors $\mathbf{E}$ and $\mathbf{B}$ will transform under the passive LT (i.e., when a basis is introduced) to another quantities, the relative vectors $\mathbf{E}'$ and $\mathbf{B}'$ *different* than $\mathbf{E}$ and $\mathbf{B}$ (the components and the basis are changed; these transformations correspond to the transformations of the usual 3D vectors $\mathbf{E}$ and $\mathbf{B}$, e.g., [11], eq. (11.149)). However the well-defined, geometric, 4D quantities, e.g., the field bivector F or *1-vectors of electric and magnetic fields*, E and B respectively, introduced below, *remain unchanged* under the passive LT (for F see above and, e.g., for E it holds that $E = E^\mu\gamma_\mu = E^{\mu'}\gamma_{\mu'}$).

These results (both with the active and the passive LT) entail that the transformations of relative vectors $\mathbf{E}$ and $\mathbf{B}$ (as parts of a well-defined 4D quantity) are not mathematically correct, which means that $\mathbf{E}$ and $\mathbf{B}$ themselves are not correctly defined quantities from the SR viewpoint. (Therefore it is not true from the SR viewpoint that ([11] Sec. 11.10): "A purely electric or magnetic field in one coordinate system will appear as a mixture of electric and magnetic fields in another coordinate frame."; or that ([3], Handout 10 in *Physical Applications of Geometric Algebra*): "Observers in relative motion see different fields.") This is very important since it shows that, in contrast to the generally accepted opinion, *the usual ME with $\mathbf{E}$ and $\mathbf{B}$ are not relativistically correct and thus they are not equivalent to the relativistically correct FE with F* (6). The same conclusion is achieved in the invariant formulation of SR with tensors [5], see particularly Sec. 5.3. in [5].

IV. THE FORMULATION OF ELECTRODYNAMICS WITH E AND B

It is clear from the above consideration that the electrodynamics cannot be correctly described from the point of view of SR with such quantities ($\mathbf{E}$ and $\mathbf{B}$) which do not transform properly under the LT.

A. Field equations with 1-vectors E and B

Therefore instead of to decompose F into $\mathbf{E}$ and $\mathbf{B}$ (23), which are not well-defined quantities from the SR viewpoint, we present *the observer independent* decomposition of F by using well-defined quantities in the Clifford algebra defined on the 4D spacetime, the vectors (grade-1) of electric E and

magnetic B fields. We define

$$F = (1/c)E \wedge v + e_5 B \cdot v. \quad (29)$$

Conversely the relations which determine E and B in terms of F are

$$E = (1/c)F \cdot v, \quad e_5 B = (1/c^2)F \wedge v, \quad B = -(1/c^2)e_5(F \wedge v), \quad (30)$$

and it holds that $E \cdot v = B \cdot v = 0$ (since F is antisymmetric), v is the velocity (1-vector) of a family of observers who measures E and B fields. The relations (29) and (30) establish the equivalence of the standard Clifford algebra formulation of electrodynamics with the field bivector F and the formulation with the 1-vectors of electric E and magnetic B fields. It is worth noting that now the observers in relative motion see the same field, e.g., the E field in the S frame is the same as in the relatively moving S' ; there is no mixture of E and B fields in S' . The FE (6) can be written in terms of 1-vectors E and B as

$$\partial((1/c)E \wedge v + e_5 B \cdot v) = j/\varepsilon_0 c. \quad (31)$$

In some basis $\{e_\mu\}$ the FE (31) can be written as

$$\begin{aligned} \partial_\alpha(\delta^{\alpha\beta}_{\mu\nu} v^\mu E^\nu - \varepsilon^{\alpha\beta\mu\nu} v_\mu c B_\nu) e_\beta &= -(j^\beta/\varepsilon_0) e_\beta, \\ \partial_\alpha(\delta^{\alpha\beta}_{\mu\nu} v^\mu c B^\nu + \varepsilon^{\alpha\beta\mu\nu} v_\mu E_\nu) e_5 e_\beta &= 0, \end{aligned} \quad (32)$$

where E^α and B^α are the basis components of the electric and magnetic 1-vectors E and B , and $\delta^{\alpha\beta}_{\mu\nu} = \delta^\alpha_\mu \delta^\beta_\nu - \delta^\alpha_\nu \delta^\beta_\mu$. The first equation in (32) (the equation with sources) emerges from $\partial \cdot F = j/\varepsilon_0 c$ and the second one (the source-free equation) is obtained from $\partial \wedge F = 0$. We remark that (32) follows from (31) for those coordinations for which the basis 1-vectors e_μ are constant, e.g., the standard basis $\{\gamma_\mu\}$ (the EC). For a nonconstant basis, for example, when one uses polar or spherical basis 1-vectors (and, e.g., the Einstein synchronization) then one must also differentiate these nonconstant basis 1-vectors. Instead of to work with F -formulation (6) one can equivalently use the E, B -formulation with the FE (31), or in the $\{e_\mu\}$ basis (32). For the given sources j one could solve these equations and find the general solutions for E and B . (We note that the equivalent formulation of electrodynamics with tensors E^a and B^a is reported in [5], while the component form in the EC is given in [8], [12] and [13].)

B. Comparison with the usual noncovariant approach with the 3-vectors $\mathbf{E}$ and $\mathbf{B}$

The comparison of this geometric approach with Clifford 1-vectors E and B and the usual noncovariant approach with the 3D vectors $\mathbf{E}$ and $\mathbf{B}$ is possible in the EC. If one considers the EC and takes that in an IFR $\mathcal{R}$ the observers who measure the basis components E^α and B^α are at rest, i.e., $v^\alpha = (c, 0, 0, 0)$, then in $\mathcal{R}$ $E^0 = B^0 = 0$. Note that we can select a particular - but otherwise arbitrary - IFR as the $\mathcal{R}$ frame, to which we shall refer as the frame of our "fiducial" observers (see [11]). In this frame of fiducial observers one can derive from the FE in the $\{\gamma_\mu\}$ basis (32) the FE which contain only the space parts E^i and B^i of E^α and B^α , e.g., from the first FE in (32) (the 1-vector part of (31)) one easily finds

$$(\partial_0 E^i - c\varepsilon^{ijk0}\partial_j B_k + j^i/c\varepsilon_0)\gamma_i - (\partial_k E^k - j^0/c\varepsilon_0)\gamma_0 = 0, \quad (33)$$

and the second FE in (32) (the trivector (pseudovector) part of (31)) yields

$$(c\partial_0 B^i + \varepsilon^{ijk0}\partial_j E_k)\gamma_5\gamma_i - (c\partial_k B^k)\gamma_5\gamma_0 = 0. \quad (34)$$

The relations (33) and (34) are coordinate-based geometric equations in the $\mathcal{R}$ frame and cannot be further simplified as geometric equations. In the equation (33) one recognizes *two ME* in the *component form*, the Ampère-Maxwell law $\varepsilon^{ijk0}\partial_j B_k = j^i/c^2\varepsilon_0 + (1/c)\partial_0 E^i$ (the first bracket, with γ_i) and the Gauss law for the electric field $\partial_i E^i = j^0/\varepsilon_0 c$ (the second bracket, with γ_0). Similarly from (34) we find the *component form* of another *two ME*, Faraday's law $\varepsilon^{ijk0}\partial_j E_k = c\partial_0 B^i$ and the Gauss law for the magnetic field $\partial_k B^k = 0$. It has to be noted that neither in $\mathcal{R}$ there is a complete mathematical equivalence of 1-vectors E and B and the 3D $\mathbf{E}$ and $\mathbf{B}$. Although the components of the 3D $\mathbf{E}$ and $\mathbf{B}$ and the components of 1-vectors E and B are the same in $\mathcal{R}$ and in the EC the usual 3D vectors and the 1-vectors, when taken as geometric quantities, i.e., together with their bases, are mathematically different quantities; the first quantities ($\mathbf{E}$ and $\mathbf{B}$) are defined on the 3D space while the second ones (E and B) are defined on the 4D spacetime. This means that from (33) and (34) one cannot derive ME with the usual 3D vectors $\mathbf{E}$ and $\mathbf{B}$, but only the component form of these equations in which E^i and B^i are the spatial components of the 1-vectors E and B (the 4D quantities). From this consideration one concludes that all the results obtained in the previous treatments from the usual ME with the 3D $\mathbf{E}$ and $\mathbf{B}$ remain valid in the formulation with the 1-vectors E and B if physical

phenomena are considered only in one IFR. Namely the selected IFR can be chosen to be the $\mathcal{R}$ frame. Then in $\mathcal{R}$, as explained above, the coordinate-based geometric equations (33) and (34) can be reduced to the equations containing only the components, the four ME in the component form. Thus for observers who are at rest in $\mathcal{R}$ ($v^\alpha = (c, 0, 0, 0)$) the components of the 3D $\mathbf{E}$ and $\mathbf{B}$ (remember that $\mathbf{E}$ and $\mathbf{B}$ are not well defined quantities from the SR viewpoint) can be simply replaced by the space components of the 1-vectors E and B in the $\{\gamma_\mu\}$ basis. It has to be noted that just such observers are usually considered in the conventional formulation with the 3D $\mathbf{E}$ and $\mathbf{B}$. However, the situation is quite different when some physical phenomena are considered from two relatively moving IFRs, say S and S' , for example, in the experiments that test SR. One of the frames, say the S frame, can be selected to be the $\mathcal{R}$ frame. Now, even in the $\mathcal{R}$ frame we cannot simply use the four ME in the component form, but from the outset we have to deal with the coordinate-based geometric equations (32), that is, (33) and (34) in $\mathcal{R}$, every of which contains two ME (the component form) together. This means that it is not correct from the SR viewpoint to investigate in two relatively moving IFRs, e.g., the Faraday law taken alone. The Faraday law is always combined with the Gauss law for the magnetic field in one unique law. They together form the relativistically correct geometric equation (32) (the second one), i.e., the equation (34) in the $\mathcal{R}$ frame. This is an important difference between our approach and all previous treatments and it will be examined in more detail in our future work. The dependence of the FE (32) on v reflects the arbitrariness in the selection of the frame $\mathcal{R}$ but at the same time it makes the equations (32) independent of that choice. The frame $\mathcal{R}$ can be selected at our disposal, which proves that we don't have a kind of the "preferred" frame theory. We see that the relativistically correct fields E and B and the new FE (31) and (32) do not have the same physical interpretation as the usual, but relativistically incorrect, 3D fields $\mathbf{E}$ and $\mathbf{B}$ and the usual 3D ME except in the frame $\mathcal{R}$ of the fiducial observers in which $E^0 = B^0 = 0$.

Furthermore an important general conclusion about the nonrelativistic physics can be drawn from the above consideration. Namely if our living arena is the 4D spacetime then only the geometric 4D quantities ($F, E, B, x, \dots$) can be correctly defined and can have an independent reality. The 3D quantities from the nonrelativistic physics, both classical and quantum, e.g., ($\mathbf{E}, \mathbf{B}, t, \mathbf{x}, E(\text{energy}), \mathbf{p}(\text{momentum}), \text{Schrödinger's } \psi, \hat{\mathbf{p}}(\text{3D momentum operator}), \dots$), do not exist by themselves, and cannot have an independent

reality in the 4D spacetime. In order that such 3D quantities can be considered as properly defined in the 4D spacetime they have to be (as in our consideration with $\mathbf{E}$ and $\mathbf{B}$) defined as parts of the corresponding geometric 4D quantities taken in a specific frame and, usually, in the EC.

C. The stress-energy vector $T(v)$ and the energy-momentum tensor $T^{\mu\nu}$ in the E, B formulation

It is shown in Sec. III that the stress-energy vector $T(v)$ for the electromagnetic field is defined in the F formulation by the relation (8). Then in the standard basis $\{\gamma_\mu\}$ we can write the stress-energy vectors T^μ as $T^\mu = T(\gamma^\mu) = (-\varepsilon_0/2)F\gamma^\mu F$. The components of the T^μ represent the energy-momentum tensor $T^{\mu\nu}$ in the $\{\gamma_\mu\}$ basis $T^{\mu\nu} = T^\mu \cdot \gamma^\nu = (-\varepsilon_0/2) \langle F\gamma^\mu F\gamma^\nu \rangle$ (remember that $\langle A \rangle$ denotes the scalar (grade-0) part of A), which reduces to familiar tensor form by writing F as $F = (1/2)F^{\mu\nu}\gamma_\mu \wedge \gamma_\nu$,

$$T^{\mu\nu} = \varepsilon_0 \left[F^{\mu\alpha} g_{\alpha\beta} F^{\beta\nu} + (1/4) F^{\alpha\beta} F_{\alpha\beta} g^{\mu\nu} \right].$$

In the usual Clifford algebra approach, e.g., [1, 2], one again makes the space-time split and considers the energy-momentum density in the γ_0 -system, $T^0 = T(\gamma^0) = T(\gamma_0)$; the split $T^0\gamma^0 = T^0\gamma_0 = T^{00} + \mathbf{T}^0$, separates T^0 into an energy density $T^{00} = T^0 \cdot \gamma^0$ and a momentum density $\mathbf{T}^0 = T^0 \wedge \gamma^0$. From the expression for T^μ and the relations (23) one finds the familiar results for the energy density $T^{00} = (\varepsilon_0/2)(\mathbf{E}^2 + c^2\mathbf{B}^2)$ and the Poynting vector $\mathbf{T}^0 = \varepsilon_0(\mathbf{E} \times c\mathbf{B})$, where the commutator product $A \times B$ is defined as $A \times B \equiv (1/2)(AB - BA)$.

However, as we have already explained, the space-time split is not relativistically correct procedure and the relative vectors $\mathbf{E}$ and $\mathbf{B}$ are not well-defined quantities on the 4D spacetime.

Therefore we express the stress-energy vector $T(v)$ in terms of 1-vectors E and B . Inserting the relation for F (29) into (10) we express $T(v)$ by means of E and B (v is again the velocity (1-vector) of a family of observers who measures E and B fields) in a compact and perspicuous form as

$$T(v) = (-\varepsilon_0/2c)(E^2 + c^2B^2)v + \varepsilon_0 e_5 [(E \wedge B) \wedge v]. \quad (35)$$

$T(v)$ defined by (35) is frame and coordination independent quantity and it is again written as a sum of the $v-\parallel$ and the $v-\perp$ parts. Thence the first term in (35) ($v-\parallel$) yields the energy density U as $U = (-\varepsilon_0/2)(E^2 + c^2B^2)$, and the second term ($v-\perp$) is $(1/c)$ of the Poynting vector $S = c\varepsilon_0 e_5 [(E \wedge B) \wedge v]$. The observer independent momentum density g is de-

defined as before $g = (1/c^2)S$ and the angular-momentum density is $M = (1/c)T(v) \wedge x = (U/c^2)v \wedge x + g \wedge x$, where $T(v)$ is given by the relation (35).

All these quantities can be written in some basis $\{e_\mu\}$ as CBGQs. Thus $T(v)$ (35) becomes

$$T(v) = (-\varepsilon_0/2c)(E^\alpha E_\alpha + c^2 B^\alpha B_\alpha)v^\lambda e_\lambda + \varepsilon_0 \tilde{\varepsilon}^\lambda_{\alpha\beta} E^\alpha B^\beta e_\lambda, \quad (36)$$

where $\tilde{\varepsilon}_{\lambda\alpha\beta} = \varepsilon_{\rho\lambda\alpha\beta}v^\rho$ is the totally skew-symmetric Levi-Civita pseudotensor induced on the hypersurface orthogonal to v . The energy density U ($U = v \cdot T(v)/c$) in the $\{e_\mu\}$ basis is determined by the first term in (36) $U = (-\varepsilon_0/2)(E^\alpha E_\alpha + c^2 B^\alpha B_\alpha)$, and the Poynting vector S in the $\{e_\mu\}$ basis is determined by the second term in (36) as $S = c\varepsilon_0 \tilde{\varepsilon}^\lambda_{\alpha\beta} E^\alpha B^\beta e_\lambda$. Of course from (36) one can easily find g and M in the $\{e_\mu\}$ basis.

Although we don't need the energy-momentum tensor $T^{\mu\nu}$ (which is defined in the $\{e_\mu\}$ basis as $T^{\mu\nu} = T^\mu \cdot e^\nu = (-\varepsilon_0/2) \langle F e^\mu F e^\nu \rangle$) we quote here $T^{\mu\nu}$ expressed in terms of components of 1-vectors E and B in some basis $\{e_\mu\}$ as

$$\begin{aligned} T^{\mu\nu} = & \varepsilon_0[(g^{\mu\nu}/2 - v^\mu v^\nu/c^2)(E^\alpha E_\alpha + c^2 B^\alpha B_\alpha) - (E^\mu E^\nu + c^2 B^\mu B^\nu) + \\ & (1/c)(\varepsilon^{\mu\alpha\beta\lambda} v_\lambda v^\nu + \varepsilon^{\nu\alpha\beta\lambda} v_\lambda v^\mu)B_\alpha E_\beta], \end{aligned} \quad (37)$$

(see also [12] and the first paper in [8] for the component form $T^{\mu\nu}$ in the EC). It has to be emphasized once again that, in contrast to all earlier definitions including the Riesz definition [14] for the energy-momentum tensor $T^{\mu\nu}$, our definitions of $T(v)$, U , S , g and M are the definitions of invariant quantities, i.e., frame and coordination independent quantities.

One can compare these expressions with familiar ones from the 3D space considering our definitions in the standard basis $\{\gamma_\mu\}$ and in the $\mathcal{R}$ frame, the frame of our fiducial observers, where $v^\alpha = (c, \mathbf{0})$, and consequently $E^0 = B^0 = 0$. Then U takes the familiar form $U = (-\varepsilon_0/2)(E^i E_i + c^2 B^i B_i)$, $i = 1, 2, 3$. Similarly, in $\mathcal{R}$, the Poynting vector becomes the familiar expression $S = \varepsilon_0 c^2 \varepsilon_0^i{}_{jk} E^j B^k \gamma_i$, $i, j, k = 1, 2, 3$, whence one also easily finds g and M in $\mathcal{R}$. Notice that all quantities in these expressions are well-defined quantities on the 4D spacetime. This again nicely illustrates our main idea that 3D quantities don't exist by themselves but only as well-defined 4D quantities taken in a particular - but otherwise arbitrary - IFR, here the $\mathcal{R}$ frame (with fiducial observers).

D. The Lorentz force in the E, B formulation

In the usual Clifford algebra approach to SR, e.g., [1 – 3], one makes the space-time split and writes the Lorentz force K in the Pauli algebra of γ_0 . Since this procedure is observer dependent we express K in an *observer independent way* using 1-vectors E and B as

$$K = (q/c) [(1/c)E \wedge v + e_5 B \cdot v] \cdot u. \quad (38)$$

In the general case when charge and observer have distinct worldlines the Lorentz force K (38) can be written as a sum of the v - $\perp$ part $K_\perp$ and the v - $\parallel$ part $K_\parallel$, $K = K_\perp + K_\parallel$, where

$$K_\perp = (q/c^2)(v \cdot u)E + (q/c)(e_5 B \cdot v) \cdot u, \quad (39)$$

$$K_\parallel = (-q/c^2)(E \cdot u)v, \quad (40)$$

respectively. This is an observer independent decomposition of the Lorentz force K . It can be easily verified that $K_\perp \cdot v = 0$ and $K_\parallel \wedge v = 0$. Both parts can be written in some basis $\{e_\mu\}$ as CBGQs $K_\perp = (q/c^2)(v^\mu u_\mu)E^\nu e_\nu + (q/c)\tilde{\varepsilon}^\mu_{\nu\rho}u^\nu B^\rho e_\mu$, where, as already said, $\tilde{\varepsilon}_{\mu\nu\rho} \equiv \varepsilon_{\lambda\mu\nu\rho}v^\lambda$, and $K_\parallel = (-q/c^2)(E^\mu u_\mu)v^\nu e_\nu$. Speaking in terms of the prerelativistic notions one can say that in the approach with the 1-vectors E and B $K_\perp$ plays the role of the usual Lorentz force lying on the 3D hypersurface orthogonal to v , while $K_\parallel$ is related to the work done by the field on the charge. However *in our invariant formulation of SR only both components together, (39) and (40), do have physical meaning and they define the Lorentz force both in the theory and in experiments.*

Let us consider a special case, the Lorentz force acting on a charge as measured by a comoving observer ($v = u$). Then from the definition of K and of E (30) one finds that $K = (q/c)F \cdot v = qE$. Thus the Lorentz force ascribed by an observer comoving with a charge is *purely electric*. The relation

$$E \equiv \lim_{q \rightarrow 0} K/q \quad (41)$$

defines the electric field E (1-vector) as the ratio of the measured force K (1-vector) on a stationary charge to the charge in the limit when the charge goes to zero. Having E so defined the charge can be given a convenient uniform velocity $u \neq v$ from which the magnetic field B (1-vector) is defined from the limit

$$[(1/c)E \wedge v + e_5 B \cdot v] \cdot u/c \equiv \lim_{q \rightarrow 0} K/q. \quad (42)$$

When *the complete* K ((38), or the sum of (39) and (40)), is known we can solve the equation of motion, Newton's second law, written as

$$(q/c) [(1/c)E \wedge v + e_5 B \cdot v] \cdot u = m(u \cdot \partial)u, \quad (43)$$

(the Lorentz force K on the l.h.s. of (43) can be also written as the sum of $K_\perp$ (39) and $K_\parallel$ (40)), $u \cdot \partial$ is the directional derivative and $(u \cdot \partial)u$ defines the acceleration of the particle. From Newton's second law (43) one obtains the force K . From K and the definitions (41) and (42) one determines the 1-vectors E and B . This form of the equation of motion differs from the usual Clifford algebra approach to SR, e.g., [1 – 3], but it agrees with the tensor formulation in general relativity, see, e.g., [15]. Notice, however, that such form (43) is used here in SR since in this invariant formulation of SR one can use different coordinations of an IFR. Thence, in general, the derivatives of the nonconstant basis vectors must be also taken into account, e.g., if one uses the Einstein synchronization and polar or spherical spatial coordinate basis. Therefore the equation of motion has to be written in the form of the equation (43).

These expressions for K can be compared with familiar ones from the 3D space considering our results in the standard basis $\{\gamma_\mu\}$ and in the $\mathcal{R}$ frame, the frame of our fiducial observers. Then $K_\perp^0 e_0 = 0$, and, e.g., $K_\perp^1 e_1 = qE^1 e_1 + q(u^2 B^3 - u^3 B^2)e_1$. Further $K_\parallel^0 e_0 = (-q/c)(E^i u_i)e_0$, and all $K_\parallel^i = 0$, $i = 1, 2, 3$. We see that in the $\mathcal{R}$ frame the whole K (38) does have $K^i = K_\perp^i$, $i = 1, 2, 3$ and, as seen from the above expressions, equal to the usual 3D expression for the Lorentz force, while $K^0 = K_\parallel^0$, and represents the usual 3D expression for the work done by the field on the charge. Then in the $\mathcal{R}$ frame the definitions of E (41) and B (42) become

$$E^i e_i \equiv \lim_{q \rightarrow 0} (K_\perp^i / q) e_i, \quad E^0 e_0 = 0 \quad (44)$$

and

$$(E^i + \varepsilon^{0kji} u_k B_j) e_i \equiv \lim_{q \rightarrow 0} (K_\perp^i / q) e_i, \quad B^0 e_0 = 0. \quad (45)$$

The expressions (44) and (45) correspond to the usual definitions of the 3D $\mathbf{E}$ and $\mathbf{B}$ in terms of the 3D expression for the Lorentz force. However it has to be noted that e_i and e_0 in (44) and (45) are the 1-vectors (defined on the 4D spacetime) while the 3D $\mathbf{E}$ and $\mathbf{B}$ and the usual 3D Lorentz force are defined on the 3D space.

V. THE FORMULATION OF ELECTRODYNAMICS WITH THE REAL $\Psi = E - ce_5 B$

In this section we consider the formulation of electrodynamics with Clifford aggregates Ψ and $\tilde{\Psi}$ and show that it is equivalent to those with F and with E and B .

$$\begin{aligned}\Psi &= E - ce_5 B, & \tilde{\Psi} &= E + ce_5 B, \\ E &= (1/2)(\Psi + \tilde{\Psi}), & B &= (1/2c)e_5(\Psi - \tilde{\Psi}).\end{aligned}\quad (46)$$

In contrast to the usual decomposition of F into the relative vectors $\mathbf{E}$ and $\mathbf{B}$ [1 – 3] *the multivectors Ψ and $\tilde{\Psi}$ defined by (46) are frame and coordination independent quantities.*

Of course it is easy to find how the F formulation and the Ψ formulation are connected

$$\begin{aligned}\Psi &= (-1/c)(vF), & \tilde{\Psi} &= (-1/c)(\tilde{F}v) = (1/c)(Fv), \\ F &= (-1/c)(v\Psi), & \tilde{F} &= (-1/c)(\tilde{\Psi}v).\end{aligned}\quad (47)$$

Then we can write the FE (6) in terms of the multivector Ψ in a simple form

$$\begin{aligned}\partial(v\Psi) &= -j/\varepsilon_0, \\ \partial \cdot (v\Psi) &= -j/\varepsilon_0, & \partial \wedge (v\Psi) &= 0.\end{aligned}\quad (48)$$

The multivector Ψ is not a homogeneous multivector but a mixed-grade multivector; it is the sum of an 1-vector and a 3-vector (pseudovector). However $v\Psi$ is a bivector (it determines F), see (47), and $v\Psi = v \cdot \Psi + v \wedge \Psi = -cv \cdot e_5 B + v \wedge E$.

The equation (48) can be written in some basis $\{e_\mu\}$ (in which the basis 1-vectors e_μ are constant) with the CBGQs

$$\partial_\alpha(\delta^{\alpha\beta}_{\mu\nu} - e_5 \varepsilon^{\alpha\beta}_{\mu\nu})v^\mu \Psi^\nu e_\beta = -(j^\beta/\varepsilon_0)e_\beta, \quad (49)$$

where $\Psi^\alpha e_\alpha = (E^\alpha - ce_5 B^\alpha)e_\alpha$. Using this last relation it can be seen that the equation (49) contains both equations with E and B from (32). The equation (49) can be also written as

$$\begin{aligned}((\Gamma^\alpha)^\beta_\mu \partial_\alpha \Psi^\mu) e_\beta &= -(j^\beta/\varepsilon_0)e_\beta, \\ (\Gamma^\alpha)^\beta_\mu &= \delta^{\alpha\beta}_{\rho\nu} v^\rho g^\nu_\mu + e_5 \varepsilon^{\alpha\beta}_{\mu\nu} v^\nu.\end{aligned}\quad (50)$$

In the case that $j^\beta = 0$ the equation (50) becomes

$$((\Gamma^\alpha)^\beta_\mu \partial_\alpha \Psi^\mu) e_\beta = 0. \quad (51)$$

It has to be emphasized that the equation (51) is a real one; Clifford algebra is developed over the field of the real numbers. Further it is not the component form in the EC but it is a coordinate-based geometric equation since it is written using CBGQs. We see from (51) that the FE for the free electromagnetic field become Dirac-like relativistic wave equation for the free photon. Really (51) is the one-photon quantum equation when Ψ is interpreted as the one-photon wave function and when the continuity equation is introduced as for the complex Ψ in [13] (the component form in the EC). However since Ψ is real there is no need for the probabilistic interpretation of Ψ ! This will be discussed elsewhere.

Let us also write the stress-energy vector $T(v)$ in terms of Ψ , and as a sum of the $v - \parallel$ and the $v - \perp$ parts. Then

$$T(v) = -(\varepsilon_0/4c) (\Psi \cdot \Psi + \tilde{\Psi} \cdot \tilde{\Psi}) v + (\varepsilon_0/4c) (\Psi \cdot \Psi - \tilde{\Psi} \cdot \tilde{\Psi}) \cdot v. \quad (52)$$

Hence $U = -(\varepsilon_0/4) (\Psi \cdot \Psi + \tilde{\Psi} \cdot \tilde{\Psi})$, and the Poynting vector is $S = (\varepsilon_0/4) (\Psi \cdot \Psi - \tilde{\Psi} \cdot \tilde{\Psi}) \cdot v$. From (52) one also finds the observer independent expressions for g and M in terms of Ψ .

All these quantities can be written in some basis $\{e_\mu\}$ as CBGQs but we will not do it here.

Only we write the energy-momentum tensor $T^{\mu\nu}$ in terms of components of Ψ in some basis $\{e_\mu\}$ as

$$\begin{aligned} T^{\mu\nu} = & \varepsilon_0 [(g^{\mu\nu}/2 - v^\mu v^\nu/c^2) \tilde{\Psi}^\alpha \Psi_\alpha - (1/2)(\tilde{\Psi}^\mu \Psi^\nu + \Psi^\mu \tilde{\Psi}^\nu) - \\ & (e_5/2c)(\varepsilon^{\mu\alpha\beta\lambda} v^\nu + \varepsilon^{\nu\alpha\beta\lambda} v^\mu) \tilde{\Psi}_\alpha \Psi_\beta v_\lambda]. \end{aligned} \quad (53)$$

The Lorentz force K is given in the real Ψ formulation as

$$\begin{aligned} K = & (q/c^2) [u \cdot (v \cdot \Psi + v \wedge \Psi)] = \\ & (q/c^2) [u \cdot (v \cdot \Psi) + (u \cdot v) \Psi - v \wedge (u \cdot \Psi)]. \end{aligned} \quad (54)$$

VI. THE FORMULATION OF ELECTRODYNAMICS WITH THE COMPLEX $\Psi = E - icB$

Sometimes it will be useful to work with the complex Ψ ; Clifford algebra is developed over the field of the complex numbers. Then

$$\begin{aligned}\Psi &= E - icB, & \bar{\Psi} &= E + icB, \\ E &= (1/2)(\Psi + \bar{\Psi}), & B &= (i/2c)(\Psi - \bar{\Psi}),\end{aligned}\tag{55}$$

i is the unit imaginary. $\bar{\Psi}$ is the complex reversion of Ψ . In contrast to the real Ψ , which is a multivector of a mixed grade (the sum of grade-1 and grade-3 multivectors), the complex Ψ is a homogeneous, grade-1, multivector. This fact facilitates the calculation in some cases, particularly when one considers the relativistic quantum mechanics. Notice also that now it holds that $v \cdot \Psi = v \cdot \bar{\Psi} = 0$.

The F formulation and the complex Ψ formulation are connected by the relations

$$\begin{aligned}\Psi &= (1/c)F \cdot v + (i/c)e_5(F \wedge v), \\ F &= (1/2c)(\Psi + \bar{\Psi}) \wedge v + (i/2c)(e_5(\Psi - \bar{\Psi})) \cdot v.\end{aligned}\tag{56}$$

Then we can use the second equation from (56) and the FE (6) to write FE in terms of the complex 1-vector Ψ as

$$\partial \cdot (v \wedge \Psi) - ie_5 [\partial \wedge (v \wedge \Psi)] = -j/\varepsilon_0.\tag{57}$$

This form of FE (in which $\bar{\Psi}$ does not appear) is achieved separating vector and trivector parts and then combining them to eliminate $\bar{\Psi}$.

Of course FE (57) can be written in some basis $\{e_\mu\}$ (with constant e_μ) with the CBGQs as

$$\partial_\alpha (\delta^{\alpha\beta}_{\mu\nu} - i\varepsilon^{\alpha\beta}_{\mu\nu}) v^\mu \Psi^\nu e_\beta = -(j^\beta/\varepsilon_0) e_\beta.\tag{58}$$

This relation is of the same form as the equation (49) but e_5 is replaced by i and in (58) Ψ is a complex 1-vector. Again it can be seen that the equation (58) contains both equations with E and B from (32). Further the equation (58) can be written in the same form as (50) only in $(\Gamma^\alpha)^\beta_\mu$ the pseudoscalar e_5 is replaced by i

$$\begin{aligned}((\Gamma^\alpha)^\beta_\mu \partial_\alpha \Psi^\mu) e_\beta &= -(j^\beta/\varepsilon_0) e_\beta, \\ (\Gamma^\alpha)^\beta_\mu &= \delta^{\alpha\beta}_{\rho\nu} v^\rho g^\nu_\mu + i\varepsilon^{\alpha\beta}_{\mu\nu} v^\nu.\end{aligned}\tag{59}$$

The equations (58) and (59) are of the same form as the corresponding equations written in the tensor formulation with the complex Ψ in [5] (or in the component form in the EC in [8] and [13]). Again for $j^\beta = 0$ we find from (59) that the FE for the free electromagnetic field become Dirac-like relativistic wave equation for the free photon.

Furthermore $T(v)$ can be expressed by Ψ and $\bar{\Psi}$ as a sum of the $v - \parallel$ and the $v - \perp$ parts, which determine U and S . Thus $T(v)$ is given as

$$T(v) = -(\varepsilon_0/2c) (\Psi \cdot \bar{\Psi}) v - i(\varepsilon_0/2c) e_5 [\Psi \wedge \bar{\Psi} \wedge v]. \quad (60)$$

Hence $U = -(\varepsilon_0/2) (\Psi \cdot \bar{\Psi})$ and $S = -i(\varepsilon_0/2) e_5 [\Psi \wedge \bar{\Psi} \wedge v]$. In some basis $\{e_\mu\}$ the stress-energy vector $T(v)$ can be written as the CBGQ

$$T(v) = (-\varepsilon_0/2c) (\Psi^\alpha \bar{\Psi}_\alpha) v^\lambda e_\lambda - i(\varepsilon_0/2c) \tilde{\varepsilon}^\lambda_{\alpha\beta} \Psi^\alpha \bar{\Psi}^\beta e_\lambda.$$

where $\tilde{\varepsilon}_{\lambda\alpha\beta} = \varepsilon_{\rho\lambda\alpha\beta} v^\rho$.

Similarly we can write the energy-momentum tensor $T^{\mu\nu}$ in terms of components of the complex Ψ in some basis $\{e_\mu\}$ and it is the same as (53) except that the pseudoscalar e_5 is replaced by i .

The Lorentz force K can be again written as a sum of the $v - \perp$ and the $v - \parallel$ parts $K = K_\perp + K_\parallel$, where

$$\begin{aligned} K_\perp &= (q/2c^2) [(u \cdot v)(\Psi + \bar{\Psi}) + i(u \wedge v \wedge (\Psi - \bar{\Psi})) e_5], \\ K_\parallel &= -(q/2c^2) [u \cdot (\Psi + \bar{\Psi}) v]. \end{aligned} \quad (61)$$

Thus all four formulations are presented in geometric terms, i.e., with invariant quantities and can be equivalently used in all calculations.

VII. COMPARISON WITH EXPERIMENTS

It is shown in [6] that the usual formulation of SR (which deals with the observer dependent quantities, i.e., the Lorentz contraction, the dilatation of time, the use of the 3D $\mathbf{E}$ and $\mathbf{B}$, etc.) shows only an "apparent" agreement (not the true one) with the traditional and modern experiments, e.g., the Michelson-Morley type experiments. On the contrary the invariant SR from [5] (given in terms of *geometric quantities - tensors*) is shown in [6] to be in a *complete agreement* with all considered experiments. This entails that the same complete agreement holds also for the formulations with *geometric quantities - the Clifford numbers*, which are presented in this paper.

In addition we briefly discuss the Trouton-Noble experiment [16] (see also [17]). In the experiment they looked for the turning motion of a charged parallel plate capacitor suspended at rest in the frame of the earth in order to measure the earth's motion through the ether. The explanations, which are given until now (see, e.g., [18]), for the null result of the experiments [16] ([17]) are not relativistically correct, since they use quantities that are not well-defined in 4D spacetime; e.g., the Lorentz contraction, the transformation equations for the usual 3D vectors $\mathbf{E}$ and $\mathbf{B}$ and for the torque as the 3D vector, the nonelectromagnetic forces of undefined nature, etc.. In our approach the explanation is very simple and natural; the energy density U , then g and M and the associated integral quantities are all invariant quantities, which means that their values are the same in the rest frame of the capacitor and in the moving frame. Thus if there is no torque (but now as a geometric, invariant, 4D quantity) in the rest frame then the capacitor cannot appear to be rotating in a uniformly moving frame.

VIII. DISCUSSION AND CONCLUSIONS

The usual Clifford algebra approach to the relativistic electrodynamics deals with the space-time split and the relative vectors $\mathbf{E}$ and $\mathbf{B}$. The investigation presented in this paper reveals that such approach is not relativistically correct. The relative vectors are not only observer dependent but their transformation law is meaningless from the SR viewpoint; it has nothing to do with the Lorentz transformations of Clifford numbers defined on the 4D spacetime. Here we employ quantities that are independent of the reference frame and of the chosen coordination for that frame. We have presented four equivalent formulations of electrodynamics by means of the field bivector F , the 1-vectors E and B , the real multivector $\Psi = E - ce_5B$ and the complex 1-vector $\Psi = E - ciB$. All four formulations are equivalent and they yield complete and consistent descriptions of electromagnetic phenomena in terms of observer independent, thus properly defined quantities on the 4D spacetime. These formulations are not equivalent with the usual Maxwell formulation with the 3D vectors $\mathbf{E}$ and $\mathbf{B}$ except in $\mathcal{R}$, the frame of fiducial observers ($v^\alpha = (c, \mathbf{0})$ and consequently $E^0 = B^0 = 0$), and when the Einstein coordination is used in $\mathcal{R}$. *The new observer independent FE* with E and B (31), with the real multivector Ψ (48) and with the complex 1-vector Ψ (57) are presented in this paper. Furthermore *the new observer independent expressions* for the stress-energy vector $T(v)$, the energy density

U , the Poynting vector S , the momentum density g , the angular-momentum density M and the Lorentz force K are given in all four formulations. The field equations with the real Ψ (51) (and the corresponding equation for the complex Ψ) for the free electromagnetic field ($j = 0$) look like Dirac relativistic wave equation for the free photon; there is no need to perform the first quantization procedure. Particularly it is important to note that in (51) the Ψ function is the real one. Hence we don't need the probabilistic interpretation for such Ψ ! The second quantization procedure, and the whole quantum electrodynamics, will be simply constructed using geometric, invariant, quantities E and B , the real or the complex Ψ , $T(v)$, U , S , g and M . Note that the standard covariant approaches to quantum electrodynamics, e.g., [19], deal with the component form (in the specific, i.e., the Einstein coordination) of the electromagnetic 4-potential A (thus requiring the gauge conditions too) instead of to use the geometric quantities, the fields 1-vectors E and B , the real or complex Ψ as Clifford numbers (this work), or the tensors E^a , B^a , or Ψ^a (see, e.g., [5]). Furthermore the standard covariant approaches employ the definitions of the field energy and momentum, which are not well-defined from the relativity viewpoint. Namely both the field energy and momentum are defined as integrals over the *three-space*, that is, over the hypersurface $t = \text{const.}$ But the hypersurface $t = \text{const.}$ in some reference frame S cannot become (under the Lorentz transformation) the hypersurface $t' = \text{const.}$ in a relatively moving reference frame S' . This is already examined for the classical electrodynamics (the covariant formulation in the EC) by Rohrlich [20] and in the first paper in [8]. Here the local conservation laws are directly derived from the FE and written in an invariant way. The observer independent integral FE and the observer independent global conservation laws (with the definitions of the invariant field energy and momentum) will be treated elsewhere. Particularly it has to be emphasized that the observer independent approach to the relativistic electrodynamics that is presented in this paper is in a complete agreement with existing experiments that test SR, which is not the case with the usual approaches. Furthermore we note that all observer independent quantities introduced here and the FE written in terms of them hold in the same form both in the flat and curved space-times. The formalism presented here will be the basis for the relativistically correct (without reference frames) formulation of quantum electrodynamics and, more generally, of the quantum field theory.

REFERENCES

- [1] D. Hestenes, *Space-Time Algebra* (Gordon and Breach, New York, 1966); *Space-Time Calculus*; available at: <http://modelingnts.la.asu.edu/evolution.html>; *New Foundations for Classical Mechanics* (Kluwer Academic Publishers, Dordrecht, 1999) 2nd. edn..
- [2] S. Gull, C. Doran, and A. Lasenby, in *Clifford (Geometric) Algebras with Applications to Physics, Mathematics, and Engineering*, W.E. Baylis, Ed. (Birkhauser, Boston, 1997), Chs. 6-8.; Found. Phys. **23**, 1175 (1993); Found. Phys. **23**, 1239 (1993); Found. Phys. **23**, 1295 (1993); Found. Phys. **23**, 1329 (1993); C. Doran, and A. Lasenby, *Physical Applications of Geometric Algebra*, available at: www.mrao.cam.ac.uk/~Clifford/
- [3] B. Jancewicz, *Multivectors and Clifford Algebra in Electrodynamics* (World Scientific, Singapore, 1989).
- [4] D. Hestenes and G. Sobczyk, *Clifford Algebra to Geometric Calculus* (Reidel, Dordrecht, 1984).
- [5] T. Ivezić, Found. Phys. **8**, 1139 (2001).
- [6] T. Ivezić, Found. Phys. Lett. **15**, 27 (2002); physics/0103026; physics/0101091.
- [7] T. Ivezić, Annales de la Fondation Louis de Broglie **27**, 287 (2002).
- [8] T. Ivezić, Found. Phys. Lett. **12**, 105 (1999); Found. Phys. Lett. **12**, 507 (1999).
- [9] A. Einstein, Ann. Physik. **17**, 891 (1905), tr. by W. Perrett and G.B. Jeffery, in *The Principle of Relativity* (Dover, New York).
- [10] D.E. Fahnline, Am. J. Phys. **50**, 818 (1982).
- [11] J.D. Jackson, *Classical Electrodynamics* (Wiley, New York, 1977) 2nd edn..
- [12] H.N. Núñez Yépez, A.L. Salas Brito, and C.A. Vargas, Revista Mexicana de Física **34**, 636 (1988).
- [13] S. Esposito, Found. Phys. **28**, 231 (1998).
- [14] M. Riesz, Dixième Congrès des Mathematicians Scandinaves, Copenhagen, 1946, p.123.
- [15] R.M. Wald, *General Relativity* (The University of Chicago Press, Chicago, 1984).
- [16] F.T. Trouton and H.R. Noble, Philos. Trans. R. Soc. London Ser. A **202**, 165 (1903).
- [17] H.C. Hayden, Rev. Sci. Instrum. **65**, 788 (1994).
- [18] A.K. Singal, J. Phys. A: Math. Gen. **25** 1605 (1992); Am. J. Phys. **61**, 428 (1993); S. A. Teukolsky, Am. J. Phys. **64**, 1104 (1996); O.D. Jefimenko, J. Phys. A: Math. Gen. **32**, 3755 (1999).

- [19] J.D. Bjorken and S.D. Drell, *Relativistic Quantum Field* (McGraw-Hill, New York, 1964); F. Mandl and G. Shaw, *Quantum Field Theory* (John Wiley & Sons, New York, 1995); S. Weinberg, *The Quantum Theory of Fields, Vol. I Foundations*, (Cambridge University Press, Cambridge, 1995).
- [20] F. Rohrlich, Phys. Rev. D **25**, 3251 (1982).